EL PASO HEALTH

ROUTINE PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (30 MONTHS - 11 YEARS)

The following guideline provides recommendations for routine preventive services for children 30 months to 11 years. Children at increased risk may warrant additional services.

The following guideline provides recommendations for routine pre Recommendation	30 mo	3 years	4 years	5 years . Cr	6 years	7 years	8 years	9 years	10 years	11 years		
History: Documentation must contain an initial health history and each				,		,			, Saile	, saile		
subsequent checkup must contain information on an interim history.	x	х	x	x	x	x	x	x	x	x		
Mental Health Screening: Mental health screening is required at each				<u> </u>				-				
check up and includes behavioral, social and emotional development.												
https://toolkits.solutions.aap.org/DocumentLibrary/BFTK2e_Links_Screeni												
ng_Tools.pdf	x	х	x	x	x	x	x	х	х	х		
Tuberculosis Screening: TB Questionnaire must be administered annually												
beginning at 12 months of age. A Tuberculin Skin Test is to be												
administered when the screening tool indicates a risk for possible exposure												
8 p		х	х	х	х	х	х	х	х	х		
Developmental Surveillance/Screening												
- Providers must use one of the following validated, standardized tools												
when performing developmental screening:												
Ages and Stages Questionnaire (ASQ)												
Ages and Stages Questionnaire: Social Emotional (ASQ:SE)												
Parents' Evaluation of Developmental Status (PEDS)												
•Survey of Well-being of young children (SWYC)												
https://toolkits.solutions.aap.org/DocumentLibrary/BFTK2e_Links_Screeni												
ng_Tools.pdf		х	х									
- Developmental surveillance is required at all other checkups and												
includes a review of milestones and mental health, including: gross and												
fine motor skills, communication skills, speech-language development, self-												
help/care skills and social, emotional, and cognitive development.	x			x	x							
Nutrition Screening: Dietary practices should be assessed to identify	^			^	^							
unusual eating habits such as pica, extended use of baby bottle feedings, or												
eating disorders in older children and adolescents. For nutritional												
problems, further assessment is indicated.	х	х	х	х	х	х	х	х	х	х		
Age Appropriate Screening & Administration of Immunizations: Providers												
must assess the immunization status of clients at every medical checkup												
and vaccines must be administered according to the current Advisory												
Committee on Immunization Practices (ACIP). "Recommended Childhood												
and Adolescent Immunization Schedule. The ACIP schedule can be found at												
https://www.cdc.gov/vaccines/schedules/	х	х	х	х	х	х	х	х	х	х		
Lead Risk Assessment Questionnaire		х	Х	Х	Х							
Laboratory Tests:												
Risk Based Test: Screenings performed based on risk assessments include												
screenings for type 2 diabetes (dia), dyslipidemia (dys), gonorrhea,												
chlamydia, syphilis (std) and HIV (hiv). Document screening or reason why										dia,dys, std,		
member was not screened.	dys	dys	dys	dys	dys	dys	dys	dys	dia, dys	hiv		
								dys (Once at 9 - 11 years)				

EL PASO HEALTH

ROUTINE PREVENTIVE SERVICES FOR CHILDREN AND ADOLESCENTS (30 MONTHS - 11 YEARS)

The following guideline provides recommendations for routine preventive services for children 30 months to 11 years. Children at increased risk may warrant additional services.

Recommendation	30 mo	3 years	4 years	5 years	6 years	7 years	8 years	9 years	10 years	11 years
Complete Unclothed Physical Examination	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
Length/Height: Requires documentation of measurements and percentiles										
as appropriate	х	х	х	х	х	х	х	х	х	х
Weight: Requires documentation of measurements and percentiles as										
appropriate	х	х	х	х	х	х	х	х	х	х
BMI: Requires documentation of measurements and percentiles as										
appropriate	х	х	х	х	х	х	х	х	х	х
Blood Pressure		х	Х	Х	Х	Х	х	Х	Х	Х

Texas Health Steps recommends that health care providers:

•Use the World Health Organization (WHO) growth charts http://www.cdc.gov/growthcharts/who_charts.htm for infants and children birth to 2 years of age.

•Use the Centers for Disease Control and Prevention (CDC) growth charts http://www.cdc.gov/growthcharts/cdc_charts.htm for children who are 2 years of age or older.

Vision: Visual Acuity results		х	х	x	х		x		x	
Subjective vision	x					x		x		x
Hearing: Audiometric Screening results			х	x	х		х		x	
Subjective hearing	х	х				х		х		х
Age Appropriate Anticipatory Guidance & Health Education: Health										
education is designed to help parents and caregivers understand what to										
expect in terms of the child's development and to provide information for										
all ages about the benefits of healthy lifestyles and practices, as well as										
accident and disease prevention. Diet and Exercise are topics that are										
strongly encouraged.										
https://hhs.texas.gov/doing-business-hhs/provider-portals/health-services-										
providers/texas-health-steps/medical-providers	Х	х	х	х	х	х	х	х	х	х
Dental : Limited oral screening for caries and general health of the teeth										
and oral mucosa is part of the physical examination. In addition to the										
federal requirements, Texas Health Steps policy requires referral to a										
dentist at six months of age and every six months thereafter until the										
dental home has been established.	Х	х	х	Х	х	Х	х	Х	Х	Х
Follow up Instructions to Return for Next Preventive Visit:	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х

This guideline is based on the Texas Health StepsPeriodicity Schedule and the Texas Health Steps Checkup Components. https://hhs.texas.gov/doing-business-hhs/provider-portals/health-services-providers/texas-health-steps/medical-providers Individual patient considerations and advances in medical science may supercede or modify these recommendations.

Revised June 1, 2021